

COVID-19 Actuarial Modelling of the Pandemic

Prepared by NMG Consulting

► Insight Report



Table of Contents

1. Multi-state actuarial model	3
1.1 Workings of the model	3
1.2 Model parameters	4
1.3 Assumptions for hospitalisations and fatalities	5
2. Setting the model parameters	5
2.1 Variable model parameters	5
2.2 Fixed model parameters	6
2.3 Viral propagation	6
3. Model fit to the South African experience	7
3.1 Calibration for the Western Cape	7
3.2 Calibration for Gauteng	8
3.3 Calibration for KwaZulu Natal	8
3.4 Calibration for the Eastern Cape	9
3.5 Calibration for the remaining provinces	10
4. Forecast results	10
4.1 Forecast of symptomatic infections	10
4.2 Forecast of hospital beds and deaths	11
4.3 Sensitivity analysis	11

NMG, along with its directors, employees, associates, affiliates, or other representatives shall not be liable for damages or injury arising out of or in connection with the use of the information contained herein including without limitation to any compensatory, direct, indirect or consequential damages, loss of data, income or profit, loss of or damage to property under any contract, negligence, strict liability or other theory arising out of or relating in any way to the information provided.

Although NMG tries to ensure that all information provided herein is correct at the time of inclusion, NMG makes no representations or warranties as to the completeness or adequacy or accuracy of the information contained herein and expressly disclaims liability for any errors or omissions or delays in updating this information. The information is supplied on the condition the persons receiving the same will make their own determination as to its suitability for their purposes prior to use or in connection with the making of any decision, and any use of same at your own risk.

1. Multi-state actuarial model

NMG has built a multi-state actuarial model to forecast COVID-19 cases in a population and applied this model to the South African and other populations. The model can be calibrated to the latest reported South Africa experience to better understand how changes in the reported experience might impact on the longer-term outlook for the pandemic.

1.1 Workings of the model

The NMG model has adopted terminology that is commonly used for modelling infectious diseases. The 'susceptible' population are those lives who are at risk of being infected. For COVID-19, the entire South African population is susceptible. Lives in the 'susceptible' population get infected by coming into contact with infectious lives. Lives so infected are allocated to the 'exposed' population while they are in the incubation stage of the disease and therefore not infectious themselves. The lives move out of the 'exposed' population to one of three infectious states after the incubation period. The three infectious states are 'symptomatic unreported' (SU), 'symptomatic reported' (SR) and 'asymptomatic unreported' (AU). From the infectious states, the lives either recover or die.

The lives needing hospitalisation and fatalities from the disease can be estimated from the lives allocated to the symptomatic infectious states and the recovered/died state.




Figure 1 Workings of the model

For each day of the disease progression, the model uses parameters to determine the likelihood of lives remaining in their current state or progressing to another state.

1.2 Model parameters

The first set of model parameters determines the progression of the disease from 'exposed' to the three infectious states as follows:

- ▶ An incubation period that determines the length of time that a life remains in the 'exposed' state before moving to one of the three infectious states;
- ▶ An infectious period that determines how long a life remains in one of the three infectious states before moving to the 'recovered/died' state; and
- ▶ A fraction that determines the proportion of lives in the infectious states that are asymptomatic.

Lives move from the 'susceptible' state to the 'exposed' state when they come into contact with a life in one of the three infectious states. The rate at which infectious lives infect those in the 'susceptible' state is determined by three factors

- ▶ A transmission rate parameter that determines the number of lives that one infectious life will infect each day;
- ▶ A relative reduction parameter applied to the transmission rate parameter for infectious lives that are asymptomatic; and
- ▶ A 'herd immunity' reduction factor calculated as the ratio of the lives in the 'susceptible' state to the lives in the overall population. This factor is very close to 1.0 at the start of the pandemic but reduces as the numbers in the recovered state increase.

1.3 Assumptions for hospitalisations and fatalities

Lives requiring hospitalisation and fatalities can be derived from the lives allocated by the model to the two asymptomatic infectious states based on a number of assumptions:

- ▶ An Infection Fatality Ratio that is used to determine the number of deaths;

- ▶ A rate of hospitalisation that is used to determine the number of potential hospital admissions;

- ▶ An average stay in hospital that is used to determine the number of potential bed days;

- ▶ An average duration from onset of the disease to hospital admission; and

- ▶ An average duration from onset of the disease to death.

2. Setting the model parameters

The NMG model parameters were initially derived from a mathematical model used to analyse the reported infections in China over the period from 10 January 2020 to 8 February 2020. Some of the model parameters have since been updated to reflect more recent information from studies on the disease progression and severity in the United Kingdom and New York.

2.1 Variable model parameters

Certain model parameters and the seed values are adjusted to fit the NMG model to the experience of the disease progression in South Africa. These variable model parameters and seed values are as follows:

- ▶ The lives allocated to the symptomatic reported and the symptomatic unreported states;

- ▶ The transmission rate parameter; and

- ▶ The Infection Fatality Ratio that is adjusted for the age distribution of the deaths due to COVID-19 in South Africa.

The adjustment to the Infection Fatality Ratio (IFR) was made using the deaths reported to 8 June 2020. This gives an average IFR that ranges from 0.468% for Gauteng to 0.743% for Eastern Cape. An adjustment has been made to the assumed IFR's to reflect an improvement of 33% to the ICU survival rates over the 30 days to 28 July 2020.

2.2 Fixed model parameters

The model parameters that are fixed in the NMG model based on international studies on the disease progression and severity are as follows:

- ▶ An incubation period of 5 days;

- ▶ An infectious period of 12 days;

- ▶ A fraction of 0.31 that determines the proportion of lives in the infectious states that are asymptomatic;

- ▶ A relative reduction parameter of 0.55 that is applied to the transmission rate parameter for infectious lives that are asymptomatic;

- ▶ A rate of hospitalisation of 5 times the deaths forecast using the Infection Fatality Ratio;

- ▶ An average stay in hospital of 7 days;

- ▶ An average duration from onset of the disease to hospital admission of 15 days; and

- ▶ An average duration from onset of the disease to death of 12 days.

The parameters for the fraction of infectious lives that are asymptomatic and the relative reduction parameter for the asymptomatic lives have been retained from the China study. There is little new evidence available that would support a change to these parameters.

The onset-to-hospital-admission parameter has been adjusted following a comparison of the reported hospitalisations and deaths for the Western Cape.

2.3 Viral propagation

The model assumes that the virus will move through a population where all individuals are equally susceptible to infection. In practice the virus will be constrained in its movements by the social contact networks that exist in the population.

The virus will infect the highly connected individuals first, and then the viral propagation will slow down when less connected individuals are left in the susceptible population.

This slowing down of the viral propagation has been built into the NMG model so as not to overstate the numbers that are forecast to eventually become infected. A parameter (% population infected parameter) must be entered that represents the percentage of the population that will have been infected when the virus stops spreading in the population.

The model transmission rate parameter will be reduced in stages until this parameter is attained to reflect this slowing down of the viral propagation.

A default ‘% population infected’ parameter has been used in the NMG model of 25% for the Western Cape and 20% for the other provinces. The default parameters for the other provinces will be increased as and when the pattern of COVID-19 deaths shows signs of exceeding the 20% level.

3. Model fit to the South African experience

The NMG model is fit (or calibrated) to the confirmed cases and deaths reported for each of the nine provinces of South Africa. The model fit is adjusted each week using the latest reported statistics.

3.1 Calibration for the Western Cape

The reported deaths in the Western Cape are compared against three scenarios for the ‘% population infected’ parameter in the graph below.




Figure 2 Forecast daily COVID-19 deaths (7-day moving average) – Western Cape

The daily reported deaths in the Western Cape spiked in the last week indicating that more than 20% of the population will be infected in the first wave of COVID-19 infections. As a result, we have increased the default ‘% population infected’ parameter to 25% for the Western Cape.

We have retained the default ‘% population infected’ parameter at 20% for the other provinces.

3.2 Calibration for Gauteng

The reported deaths in Gauteng are compared against three scenarios for the ‘% population infected’ parameter in the graph below.




Figure 3 Forecast daily COVID-19 deaths (7-day moving average) – Gauteng

The experience in Gauteng continues to track below the 15% population infected level and the forecast for this province will be recalibrated in our next report if the trend continues.

3.3 Calibration for KwaZulu Natal

The reported deaths in KwaZulu Natal are compared against three scenarios for the ‘% population infected’ parameter in the graph below.




Figure 4 Forecast daily COVID-19 deaths (7-day moving average) – KwaZulu Natal

The experience in KwaZulu Natal continues to track below the 15% population infected level and the forecast for this province will be recalibrated in our next report if the trend continues.

3.4 Calibration for the Eastern Cape

The reported deaths in the Eastern Cape are compared against three scenarios for the ‘% population infected’ parameter in the graph below. There was a delay in the reporting of deaths that has impacted the model fit.




Figure 5 Forecast daily COVID-19 deaths (7-day moving average) – Eastern Cape

3.5 Calibration for the remaining provinces

The shape of the first wave derived from Gauteng and KwaZulu Natal is applied to the remaining provinces to get a model fit for the country as a whole.

4. Forecast results

NMG has released a web-based version of the multi-state actuarial model that will allow interested parties to generate different COVID-19 forecast scenarios for South Africa. The model can be accessed at <https://nmg-covid-19.sctechnology.co.za/>

4.1 Forecast of symptomatic infections

The symptomatic infections implied by the NMG model are plotted for the four large provinces in the table below using the default ‘% population infected’ parameters. Only for the Western Cape, do the infections reach a plateau. The model will be adjusted if a similar plateau shape becomes evident in any of the other provinces.




Figure 6 COVID-19 symptomatic infections projected for top four provinces

Taking an aggregate of all nine provinces provides a forecast for the aggregate of symptomatic and asymptomatic infections that shows them having already peaked. The forecast of infections for South Africa as whole is shown below:




Figure 7 Forecast of reported and unreported infections

4.2 Forecast of hospital beds and deaths

The number of potential lives being treated in hospital for COVID-19 is forecast to start coming off its peak at 12 600. Based on adjusted international Infection Fatality Ratios applied at a province level and the default '% population infected' parameters, some 31 500 lives are expected to be lost due to the pandemic.




Figure 8 Forecast of hospital beds and cumulative deaths

4.3 Sensitivity analysis

The graphs below expand on how the input parameters impact of the model projection results.

The ‘% population infected’ parameter is key to determining the number of lives in the population that will become infected by the virus in this first wave. The projected symptomatic infections are shown below for the default parameter value and values 5% less and 5% more than the default:




Figure 9 Forecast of COVID-19 infections (symptomatic)

The forecast cumulative COVID-19 deaths are shown below for the default parameter value for ‘% population infected’ and values 5% less and 5% more than the default:




Figure 10 Forecast of cumulative COVID-19 deaths

The forecast for hospital bed days required for COVID-19 patients are shown below for the default parameter value for ‘% population infected’ and values 5% less and 5% more than the default:




Figure 11 Forecast of potential hospital beds required due to COVID-19

Thank you

For more information,
visit www.nmg-consulting.com